

Patient Name	: Mr.Dummy	Collected	: 08/Jul/2026 09:48AM
Age/Gender	: 67 Y 0 M 0 D /M	Received	: 08/Jul/2026 10:26AM
UHID/MR No	: Dummy.0000000	Reported	: 08/Jul/2026 10:55AM
Visit ID	: Dummy101	Status	: Final Report
Ref Doctor	: Dr.SELF	Client Name	: NRL HYDERABAD LAB
IP/OP NO	:	Patient location	: Dummy Location

DEPARTMENT OF BIOCHEMISTRY

Test Name	Result	Unit	Bio. Ref. Interval	Method
-----------	--------	------	--------------------	--------

C-REACTIVE PROTEIN CRP (QUANTITATIVE) , <i>SERUM</i>	70.37	mg/L	0-5	Latex Particle Immunoturbidimetric
---	--------------	------	-----	------------------------------------

Comment:

C-reactive protein (CRP) is one of the most sensitive acute-phase reactants for inflammation. Measuring changes in the concentration of CRP provides useful diagnostic information about the level of acuity and severity of a disease. Unlike ESR, CRP levels are not influenced by hematologic conditions such as anemia, polycythemia etc.

Increased levels are consistent with an acute inflammatory process. After onset of an acute phase response, the serum CRP concentration rises rapidly (within 6-12 hours and peaks at 24-48 hours) and extensively. Concentrations above 100 mg/L are associated with severe stimuli such as major trauma and severe infection (sepsis).

***** End Of Report *****

