Patient Name	: Mr.Dummy	Collected	: 08/Jul/2026 10:02AM
Age/Gender	: 37 Y 10 M 28 D/M	Received	: 08/Jul/2026 10:29AM
UHID/MR No	: Dummy.0000000	Reported	: 08/Jul/2026 12:03PM
Visit ID	: Dummy101	Status	: Final Report
Ref Doctor	: Dr.SELF	Client Name	: NRL HYDERABAD LAB
IP/OP NO	:	Patient location	: Dummy Location

DEPARTMENT OF IMMUNOLOGY

Test Name	Result	Unit	Bio. Ref. Interval	Method
FREE T4 (FT4) , *SERUM*	1.00	ng/dL	0.8 - 2.7	CMIA

Comment:

FT4 gives corrected values in patients in whom the total T4 is altered on account of changes in serum proteins or in binding sites. Monitoring restoration to normal range is the only laboratory criterion to estimate appropriate replacement dose of levothyroxine because 6–8 weeks are required before TSH reflects these changes

FT4 assays are prone to inaccurate readings in pregnant women. Anticonvulsant drug therapy (particularly phenytoin) may result in decreased FT4 levels due to an increased hepatic metabolism and secondarily to displacement of hormone from binding sites.

***** End Of Report *****

SIN No:Dummyb001

This test has been performed at Apollo Health & Lifestyle Ltd,Diagnostic Lab,Bowenpally,Secunderabad